

## Canadian Community Health Survey Statistics – Canada

|                                     | 2001              | 2002* | 2003              | 2004* | 2005              | 2006* | 2007** | 2008** | 2009** | 2010              | 2011** | % change<br>2005-2010 |
|-------------------------------------|-------------------|-------|-------------------|-------|-------------------|-------|--------|--------|--------|-------------------|--------|-----------------------|
| <b>Frequency Total</b>              | <b>25,787,334</b> |       | <b>26,555,430</b> |       | <b>27,126,165</b> |       |        |        |        | <b>28,878,418</b> |        | <b>6%</b>             |
| Chronic Fatigue Syndrome            | 203,280           |       | 340,307           |       | 333,816           |       |        |        |        | 411,465           |        | 23%                   |
| Fibromyalgia                        | 279,751           |       | 392,492           |       | 389,782           |       |        |        |        | 438,831           |        | 13%                   |
| Multiple Chemical Sensitivities     | 475,436           |       | 642,808           |       | 598,585           |       |        |        |        | 800,333           |        | 34%                   |
| None of the three conditions        | 24,951,787        |       | 25,372,033        |       | 25,990,940        |       |        |        |        | 27,463,696        |        | 6%                    |
| One or more of the three conditions | 835,547           |       | 1,183,397         |       | 1,135,225         |       |        |        |        | 1,414,722         |        | 25%                   |
| <b>Prevalence Total</b>             |                   |       |                   |       |                   |       |        |        |        |                   |        |                       |
| Chronic Fatigue Syndrome            | 0.8%              |       | 1.3%              |       | 1.2%              |       |        |        |        | 1.4%              |        |                       |
| Fibromyalgia                        | 1.1%              |       | 1.5%              |       | 1.4%              |       |        |        |        | 1.5%              |        |                       |
| Multiple Chemical Sensitivities     | 1.8%              |       | 2.4%              |       | 2.2%              |       |        |        |        | 2.8%              |        |                       |
| None of the three conditions        | 96.8%             |       | 95.5%             |       | 95.8%             |       |        |        |        | 95.1%             |        |                       |
| One or more of the three conditions | 3.2%              |       | 4.5%              |       | 4.2%              |       |        |        |        | 4.9%              |        |                       |

Extracted from the CCHS Public Use Microdata Files (unrounded)  
by Margaret Parlor

31/01/2013

\* The Canadian Community Health Survey was not run in 2002, 2004 or 2006

\*\* The questions on CFS, FM and MCS were not included on the Survey in 2007, 2008, 2009 or 2011.